

HAMILTON, BROOK, SMITH & REYNOLDS, P.C.

06-09-00

PTO
JCS30 U.S.
09/590211



A

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under
37 C.F.R. 1.53(b))

Attorney Docket No.

3028.1000-000

First Named Inventor or
Application Identifier

Guy A. Rouleau

Express Mail Label No.

EL564265605US

Title of
Invention

SHORT GCG EXPANSIONS IN THE PAB II GENE FOR OCULOPHARYNGEAL
MUSCULAR DYSTROPHY AND DIAGNOSTIC THEREOF

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Assistant Commissioner for Patents
Box Patent Application
Washington, D.C. 20231

1. ☐ Fee Transmittal Form
(Submit an original, and a duplicate for fee processing)

2. ☒ Specification **Total Pages 20**
(preferred arrangement set forth below)

- Descriptive title of the invention
- Cross References to Related Applications
- Statement Regarding Fed sponsored R & D
- Reference to microfiche Appendix
- Background of the Invention
- Summary of the Invent on
- Brief Description of the Drawings
- Detailed Description
- Claim(s)
- Abstract of the Disclosure

3. ☒ Drawing(s) (35 U.S.C. 113) **Total Sheets 8**
☐ Formal ☒ Informal

4. ☐ Oath or Declaration/POA **[Total Pages []]**

- a. ☐ Newly executed (original or copy)
- b. ☐ Copy from a prior application (37 C.F.R. 1.63(d))
(for continuation/divisional with Box 17 completed)
[NOTE Box 5 below]
 - i. ☐ DELETION OF INVENTOR(S)
Signed statement attached deleting
inventor(s) named in the prior
application, see 37 C.F.R. 1.63(d)(2)
and 1.33(b).

5. ☐ Incorporation By Reference (useable if Box 4b is checked)
The entire disclosure of the prior application, from which a
copy of the oath or declaration is supplied under Box 4b, is
considered as being part of the disclosure of the accompanying
application and is hereby incorporated by reference therein.

6. ☐ Microfiche Computer Program (Appendix)

7. ☒ Nucleotide and/or Amino Acid Sequence Submission
(if applicable, all necessary)

- a. ☐ Computer Readable Copy
- b. ☒ Paper Copy (identical to computer copy)
3 Pages (1/3-3/3)
- c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet & documents)

9. ☐ 37 C.F.R. 3.73(b) Statement ☐ Power of Attorney
(when there is an assignee)

10. ☐ English Translation Document (if applicable)

11. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations

12. ☐ Preliminary Amendment

13. ☒ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)

14. ☐ Small Entity Statement(s) ☐ Statement filed in prior application,
status still proper and desired

15. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)

16. ☒ Other: Remarks Letter

17. If a **CONTINUING APPLICATION**, check appropriate box and supply the requisite information:

☒ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.: PCT/CA98/01133

Prior application information: Examiner:

Group Art Unit:

18. CORRESPONDENCE ADDRESS

NAME

David E. Brook, Esq.

HAMILTON, BROOK, SMITH & REYNOLDS, P.C.

ADDRESS

Two Militia Drive

CITY

Lexington

STATE

MA

ZIP CODE

02421-4799

COUNTRY

USA

TELEPHONE

(781) 861-6240

FAX

(781) 861-9540

Signature

David E. Brook, R.N. 22592 for
Elizabeth W. Mata

Date

6/8/00

Submitted by
Typed or Printed Name

Elizabeth W. Mata

Reg. Number

38,236

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants: Guy A. Rouleau and Bernard Brais

Continuation Application of International

Application No.: PCT/CA98/01133

Filed: December 7, 1998

For: SHORT GCG EXPANSIONS IN THE PAB II GENE FOR OCULOPHARYNGEAL
MUSCULAR DYSTROPHY AND DIAGNOSTIC THEREOF

Date: June 8, 2000

EXPRESS MAIL LABEL NO. EL56426560SL5

REMARKS

Box PATENT APPLICATION

Assistant Commissioner for Patents

Washington, D.C. 20231

Sir:

The referenced application is a Continuation application claiming priority to PCT International Application No. PCT/CA98/01133, filed 7 December 1998, which claimed priority to Canadian Patent No. 2,218,199.

The application has been amended as follows from the original PCT filing: a Related Applications paragraph has been added; the description of Fig. 4 has been changed to reflect the actual labeling of the figure as Figs. 4A-4E; the specification has been amended to correct a number of typographical and syntax mistakes or irregularities; information previously residing in the Background of the Invention has been transferred to the Summary of the Invention, where it properly belongs; a description of a reference cited in the International Search report (Brais *et al.*) has been added to the Description of the Prior Art in the Background of the Invention; proper units have been inserted (e.g., " μ ") in the description of "PAB II mutation screening and

sequencing" in the Examples; and trademarks have been appropriately identified. No new matter has been added.

The claims in the Continuation application are based on the set of claims submitted with the Reply to the First Written Opinion with regard to the original PCT filing. The claims have been amended from those in the Reply to the First Written Opinion, in that multiple dependencies have been deleted, and the gene of Claims 1 and 31 has been specified to be an "isolated" gene.

To facilitate analysis of the claims and to emphasize that no new matter has been added, representative support in the original PCT filing is identified herein for the claims in the Continuation application. All support references pertain to pagination in the original PCT filing.

Support for Claims 1-12 and 31-32 can be found, for example, in Figure 2, at page 2, lines 12-1, at page 6, line 9, through page 7, line 6, and page 7, line 33 through page 8, line 19, and page 8, lines 26-31. Additional support for Claim 1 can also be found, for example at page 1, line 15; support for Claim 2, see Figure 2 and page 5, lines 24-30; support for Claims 5 and 6, see page 8, lines 11-12; support for Claims 7 and 8, see page 6, lines 9-17.

Support for Claim 13, can be found, for example, in Claim 6, at page 2, lines 17-31; support for Claims 14-18, which depend from Claim 13, can be found, for example, at page 7, line 33 through page 8, line 31. Support for Claims 19-24 can be found, for example, in old Claim 9, and at page 2, line 32, through page 3, line 3; support for Claims 25-27, see old Claim 10; support for Claim 28-30, see page 2, line 32, through page 3, line 3; support for Claim 33-36, see at page 7, line 33 through page 8, line 31.

If the Examiner believes that a telephone call would expedite prosecution of the application, the Examiner is invited to contact Elizabeth W. Mata at (915) 845-3558. If Elizabeth W. Mata cannot be reached, the Examiner is invited to contact David E. Brook at (781) 861-6240.

Respectfully submitted,

HAMILTON, BROOK, SMITH & REYNOLDS, P.C.

David E. Brook RN 22592

By *for Elizabeth W. Mata*

Elizabeth W. Mata

Registration No. 38,236

Telephone: (915) 845-3558

Facsimile: (915) 845-3237

Lexington, Massachusetts 02421-4799

Dated: 6/8/00

-1-

Date: <u>June 8, 2000</u>	Express Mail Label No. <u>EL56426560SL5</u>
---------------------------	---

Inventors: Guy A. Rouleau and Bernard Brais

Attorney's Docket No.: 3028.1000-000

SHORT GCG EXPANSIONS IN THE PAB II GENE FOR OCULO-
PHARYNGEAL MUSCULAR DYSTROPHY AND DIAGNOSTIC THEREOF

RELATED APPLICATION(S)

This application is a continuation claiming priority to International Application
5 No. PCT/CA98/01133, filed December 7, 1998 (designating the U.S.), which claims
priority to Canadian Patent No. 2,218,199, filed December 9, 1997, the entire teachings
of both of which are incorporated herein by reference.

BACKGROUND OF THE INVENTION

Field of the Invention

10 The invention relates to PAB II gene, and its uses thereof for the diagnosis,
prognosis and treatment of a disease related with protein accumulation in nucleus, such
as oculopharyngeal muscular dystrophy.

Description of Prior Art

Autosomal dominant oculopharyngeal muscular dys-rophy(OPMD) is an adult-
15 onset disease with a world-wide distribution. It usually presents itself in the sixth
decade with progressive swallowing difficulties (dys-phagia), eye lid drooping (ptosis)
and proximal limb weakness. Unique nuclear filament inclusions in skeletal muscle
fibers are its pathological hallmark (Tome, F.M.S. & Fardeau, Acta Neuropath. 49, 85-
87 (1980)). Using the full power of linkage analysis in eleven French Canadian families,

the oculopharyngeal muscular dystrophy gene was fine mapped on human chromosome 14 (Brais et al., 1997, Neuromuscular Disorders 7 (Suppl.1):S70-74). A region of .75 cM was thereby identified as a region containing the potential and unknown OMPD gene (Brais et al., 1997, *supra*). Unfortunately, the OMPD gene has yet to be isolated and its nucleic acid or protein sequence have yet to be cribbed.

It would be highly desirable to be provided with a tool for the diagnosis, prognosis and treatment of a disease related with polyalanine accumulation in the nucleus, such as observed in oculopharyngeal muscular dystrophy.

SUMMARY OF THE INVENTION

One aim of the present invention is to provide a tool for the diagnosis, prognosis and treatment of a disease related with polyalanine accumulation in nucleus, such as oculopharyngeal muscular dystrophy.

Herein, the poly(A) binding protein II (PAB II) gene was isolated from a 217 kb candidate interval in chromosome 14q11.A (GCG)6 repeat encoding a polyalanine tract located at the N-terminus of the protein was expanded to (GCG)8-13 in the 144 OPMD families screened. More severe pheno-types were observed in compound heterozygotes for the (GCG)9 mutation and a (GCG)7 allele found in 2% of the population, whereas homozygosity for the (GCG)7 allele leads to autosomal recessive OPMD. Thus the (GCG)7 allele is an example of a polymorphism which can act as either a modifier of a dominant phenotype or as a recessive mutation. Pathological expansions of the polyalanine tract may cause mutated PAB II oligomers to accumulate as filament inclusions in nuclei.

In accordance with the present invention there is provided a human PAB II gene containing a transcribed polymorphic GCG repeat, which comprises a sequence as set forth in Fig. 4, which includes introns and flank-ing genomic sequence.

The allelic variants of GCG repeat of the human PAB II gene are associated with a disease related with protein accumulation in the nucleus, such as polyalanine

accumulation, or with a disease related with swallowing difficulties, such as oculopharyngeal muscular dystrophy.

In accordance with the present invention there is also provided a method for the diagnosis of a dis-ease associated with protein accumulation in the nucleus, which

5 comprises the steps of:

- a) obtaining a nucleic acid sample of said patient; and
- b) determining allelic variants of a GCG repeat of the human PAB II gene; thereby long allelic variants are indicative of a disease related with protein accumulation in the nucleus, such as polyalanine accumulation and oculopharyngeal muscular dystrophy.

10

The long allelic variants have from about 245 to about 263 bp in length.

In accordance with the present invention there is also provided a non-human mammal model for the human PAB II gene, whose germ cells and somatic cells are modified to express at least one allelic variant of the PAB II gene and wherein said

15 allelic variant of the PAB II is being introduced into the mammal, or an ancestor of the mammal, at an embryonic stage.

In accordance with the present invention there is also provided a method for the screening of therapeutic agents for the prevention and/or treatment of oculopharyngeal muscular dystrophy, which comprises the steps of:

- 20 a) administering the therapeutic agents to the non-human animal of the present invention or oculopharyngeal muscular dystrophy patients; and
- b) evaluating the prevention and/or treatment of development of oculopharyngeal muscular dystrophy in this animal (such as a mammal) or in patients.

In accordance with the present invention there is also provided a method to
25 identify genes-products thereof, or part thereof, which interact with a biochemical pathway affected by the PAB II gene, which comprises the steps of:

- a) designing probes and/or primers using the PAB II gene and screening oculopharyngeal muscular dystrophy patients samples with said probes and/or primers, and
- b) evaluating the role of the identified gene in oculopharyngeal muscular dystrophy patients.

BRIEF DESCRIPTION OF THE DRAWINGS

Figs. 1A-1B illustrate the positional cloning of the PAB II gene;

Figs. 2A-2G illustrate the OPMD (GCG)_n expansion sizes and sequence of the mutation site (SEQ ID NOS:1-2);

Fig. 3 illustrates the age distribution of swallowing time (st) for French Canadian OPMD carriers of the (GCG)₉ mutation; and

Figs. 4A-4E illustrate the nucleotide sequence of human poly(A) binding protein II (hPAB II)(SEQ ID NO:3).

DETAILED DESCRIPTION OF THE INVENTION

In order to identify the gene mutated in OPMD, a 350 kb cosmid contig was constructed between flanking markers D14S990 and D14S1457 (Fig. 1A). Positions of the PAB II-selected cDNA clones were determined in relation to the EcoRI restriction map and the Genealogy-based Estimate of Historical Meiosis (GEHM)-derived candidate interval (Rommens, J.M. et al., in Proceedings of the third international workshop on the identification of transcribed sequences (eds. Hochgeschwender, U. & Gardiner, K.) 65-79 (Plenum, New York, 1994)).

The human poly(A) binding protein II gene (PAB II) is encoded by the nucleotide sequence as set forth in Fig. 4.

Twenty-five cDNAs were isolated by cDNA selection from the candidate interval (Rommens, J.M. et al., in Proceedings of the third international workshop on the identification of transcribed sequences (eds. Hochgeschwender, U. & Gardiner, K.;

65-79; Plenum, New York, 1994). Three of these hybridized to a common 20 kb EcoRI restriction fragment and showed high sequence homology to the bovine poly(A) binding protein II gene (bPAB II) (Fig. 1A). The PAB II gene appeared to be a good candidate for OPMD because it mapped to the genetically defined 0.26 cM candidate interval in 14q11 (Fig. 1A), its mRNA showed a high level of expression in skeletal muscle, and the PAB II protein is exclusively localized to the nucleus (Krause, S. et al., Exp. Cell Res. 214, 75-82 (1994)) where it acts as a factor in mRNA polyadenylation (Whale, E., Cell 66, 759-768 (1991); Whale, E. et al., J. Biol. Chem. 268, 2937-2945 (1993); Bienroth, S. et al., EMBO J. 12, 585-594 (1993)).

10 A 8 kb HindIII genomic fragment containing the PAB II gene was subcloned and sequenced (6002 bp; GenBank: AF026029)(Nemeth, A. et al., Nucleic Acids Res. 23, 4034-4041 (1995)) (Fig. 1B). Genomic structure of the PAB II gene, and position of the OPMD (GCG)_n expansions. Exons are numbered. Introns 1 and 6 are variably present in 60% of cDNA clones. ORF, open reading frame; cen, centromere and tel, 15 telomere.

The coding sequence was based on the previously published bovine sequence (GenBank: X89969) and the sequence of 31 human cDNAs and ESTs. The gene is composed of 7 exons and is transcribed in the centromere orientation (Fig. 1B). Multiple splice variants are found in ESTs and on Northern blots (Nemeth, A. et al., Nucleic 20 Acids Res. 23, 4034-4041 (1995)). In particular, introns 1 and 6 are present in more than 60% of clones (Fig. 1B)(Nemeth, A. et al., Nucleic Acids Res. 23, 4034-4041 (1995)). The coding and protein sequences are highly conserved between human, bovine and mouse (GenBank: U93050). 93% of the PAB II sequence was readily amenable to RT-PCR- or genomic-SSCP screening. No mutations were uncovered 25 using both techniques. However, a 400 bp region of exon 1 containing the start codon could not be readily amplified. This region is 80% GC rich. It includes a (GCG)₆ repeat which codes for the first six alanines of a homopolymeric stretch of 10 (Fig. 2G). Nucleotide sequence of the mutated region of PAB II as well as the amino acid

sequences of the N-terminus polyalanine stretch and position of the OPMD alanine insertions is also shown in Fig. 2.

Special conditions were designed to amplify by PCR a 242 bp genomic fragment including this GCG-repeat. The (GCG)6 allele was found in 98% of French Canadian non-OPMD control chromosomes, whereas 2% of chromosomes carried a (GCG)7 polymorphism (n=86) (Brais, B. et al., Hum. Mol. Genet. 4, 429-434 (1995)).

Screening OPMD cases belonging to 144 families showed in all cases a PCR product larger by 6 to 21 bp than that found in controls (Fig. 2A). (GCG)6 normal allele (N) and the six different (GCG)n expansions observed in 144 families.

Sequencing of these fragments revealed that the increased sizes were due to expansions of the GCG repeat (Fig. 2G). Fig. 2F shows the sequence of the (GCG)9 French Canadian expansion in a heterozygous par-ent and his homozygous child. Partial sequence of exon 1 in a normal (GCG)6 control (N), a heterozygote (ht.) and a homozygote (hrn.) for the (GCG)9-repeat mutation. The number of families sharing the different (GCG)n-repeats expansions is shown in Table 1.

Table 1
Number of families sharing the different dominant (GCG)n OPMD mutations

5	Mutations	Polyalanine†	Families
	(GCG)8	12	4
	(GCG)9	13	99
	(GCG)10	14	19
10	(GCG)11	15	16
	(GCG)12	16	5
	(GCG)13	17	1
	Total		144

†, 10 alanine residues in normal PAB II.

15 The (GCG)9 expansion shared by 70 French Canadian families is the most frequent mutation we observed (Table 1).The (GCG)9 expansion is quite stable, with a single doubling observed in family F151 in an estimated 598 French Canadian meioses (Fig. 2C). The doubling of the French Canadian (GCG)9 expansion is demonstrated in Family F151.

20 This contrasts with the unstable nature of previously described disease-causing triplet-repeats (Rosenberg, R.N., New Eng. J. Med. 335, 1222-1224 (1996)).

 Genotyping of all the participants in the clinical study of French Canadian OPMD provided molecular insights into the clinical variability observed in this condition. The genotypes for both copies of the PAB II mutated region were added to an

25 anonymous version of this clinical database of 176 (GCG)9 mutation carriers (Brais, B. et al., Hum. Mol. Genet. 4, 429-434 (1995)). Severity of the phenotype can be assessed by the swallowing time (st) in seconds taken to drink 80 cc of ice-cold water (Brais, B.

et al., Hum. Mol. Genet. 4, 429-434 (1995); Bouchard, J.-P. et al., Can. J. Neurol. Sci. 19, 296-297 (1992)). The late onset and progressive nature of the muscular dystrophy is clearly illustrated in heterozygous carriers of the (GCG)9 mutation (bold curve in Fig. 3) when compared to the average st of control (GCG)6 homozygous

- 5 participants ($n=76$, thinner line in Fig. 3). The bold curve represents the average OPMD st for carriers of only one copy of the (GCG)9 mutation ($n=169$), while the thinner line corresponds to the average st for (GCG)6 homozygous normal controls ($n=76$). The black dot corresponds to the st value for individual VIII. Roman numerals refer to individual cases shown in Figs. 2B, 2D and discussed in the text. The genotype of a
- 10 homozygous (GCG)9 patient and her parents is shown in Fig. 2B. Independent segregation of the (GCG)7 allele is also shown. Of note, case V has a more severe OPMD phenotype (Fig. 2D).

- Two groups of genotypically distinct OPMD cases have more severe swallowing difficulties. Individuals I, II, and III have an early-onset disease and are
- 15 homozygous for the (GCG)9 expansion ($P < 10^{-5}$) (Figs. 2B, F). Cases IV, V, VI and VII have more severe phenotypes and are compound heterozygotes for the (GCG)9 mutation and the (GCG)7 polymorphism ($P < 10^{-5}$). In Fig. 2D the independent segregation of the two alleles is shown. Case V, who inherited the French Canadian (GCG)9 mutation and the (GCG)7 polymorphism, is more symptomatic than his brother
 - 20 VIII who carries the (GCG)9 mutation and a normal (GCG)6 allele (Figs. 2D and 3). The (GCG)7 polymorphism thus appears to be a modifier of severity of dominant OPMD. Furthermore, the (GCG)7 allele can act as a recessive mutation. This was documented in the French patient IX who inherited two copies of the (GCG)7 polymorphism and has a late-onset autosomal recessive form of OPMD (Fig. 2E). Case
 - 25 IX, who has a recessive form of OPMD, is shown to have inherited two copies of the (GCG)7 polymorphism.

This is the first description of short trinucleotide repeat expansions causing a human disease. The addition of only two GCG repeats is sufficient to cause dominant

OPMD. OPMD expansions do not share the cardinal features of "dynamic mutations". The GCG expansions are not only short they are also meiotically quite stable. Furthermore, there is a clear cut-off between the normal and abnormal alleles, a single GCG expansion causing a recessive phenotype. The PAB II (GCG)⁷ allele is the first

5 example of a relatively frequent allele which can act as either a modifier of a dominant phenotype or as a recessive mutation. This dosage effect is reminiscent of the one observed in a homozygote for two dominant synpolydactyly mutations. In this case, the patient had more severe deformities because she inherited two duplications causing an expansion in the polyalanine tract of the HOXD13 protein (Akarsu, A.N. et al., Hum.

10 Mol. Genet. 5, 945-952 (1996)). A duplication causing a similar polyalanine expansion in the a subunit 1 gene of the core-binding transcription factor (CBF(1) has also been found to cause dominant cleido-cranial dysplasia (Mundlos, S. et al., Cell 89, 773-779 (1997)). The mutations in these two rare diseases are not triplet-repeats. They are duplications of "cryptic repeats" composed of mixed synonymous codons and are

15 thought to result from unequal crossing over (Warren, S.T., Science 275, 408-409 (1997)). In the case of OPMD, slippage during replication causing a reiteration of the GCG codon is a more likely mechanism (Wells, D.R., J. Biol. Chem. 271, 2875-2878 (1996)).

Different observations converge to suggest that a gain of function of PAB II

20 may cause the accumulation of nuclear filaments observed in OPMD (Tome, F.M.S. & Fardeau, Acta Neuropath. 49, 85-87 (1980)). PAB II is found mostly in dimeric and oligomeric forms (Nemeth, A. et al., Nucleic Acids Res. 23, 4034-4041 (1995)). It is possible that the polyalanine tract plays a role in polymerization. Polyalanine stretches have been found in many other nuclear proteins such as the HOX proteins, but their

25 function is still unknown (Davies, S.W. et al., Cell 90, 537-548 (1997)). Alanine is a highly hydrophobic amino acid present in the cores of proteins. In dragline spider silk, polyalanine stretches are thought to form β -sheet structures important in ensuring the fibers' strength (Simmons, A.H. et al., Science 271, 84-87 (1996)). Polyalanine

oligomers have also been shown to be extremely resistant to chemical denaturation and enzymatic degradation (Forood, B. et al., *Bioch. and Biophys. Res. Com.* 211, 7-13 (1995)). One can speculate that PAB II oligomers comprised of a sufficient number of mutated molecules might accumulate in the nuclei by forming undegradable polyaniline rich macromolecules. The rate of the accumulation would then depend on the ratio of mutated to non-mutated protein. The more severe phenotypes observed in homozygotes for the (GCG)₉ mutations and compound heterozygotes for the (GCG)₉ mutation and (GCG)₇ allele may correspond to the fact that in these cases PAB II oligomers are composed only of mutated proteins. The ensuing faster filament accumulation could cause accelerated cell death. The recent description of nuclear filament inclusions in Huntington's disease, raises the possibility that "nuclear toxicity" caused by the accumulation of mutated homopolymeric domains is involved in the molecular pathophysiology of other triplet-repeat diseases (Davies, S.W. et al., *Cell* 90, 537-548 (1997); Scherzinger, E. et al., *Cell* 90, 549-558 (1997); DiFiglia, M. et al., *Science* 277, 1990-1993 (1997)). Future immunocytochemical and expression studies will be able to test this patho-physiological hypothesis and provide some insight into why certain muscle groups are more affected while all tissues express PAB II.

Methods

Contig and cDNA selection

The cosmid contig was constructed by standard cosmid walking techniques using a gridded chromosome 14-specific cosmid library (Evans, G.A. et al., *Gene* 79, 9-20 (1989)). The cDNA clones were isolated by cDNA selection as previously described (Rommens, J.M. et al., in *Proceedings of the third international workshop on the identification of transcribed sequences* (eds. Hochgeschwender, U. & Gardiner, K.) 65-79 (Plenum, New York, 1994)).

Cloning of the PAB II gene

Three cDNA clones corresponding to PAB II were sequenced (Sequenase, USB). Clones were verified to map to cosmids by South-ern hybridization. The 8 kb HindIII restriction frag-ment was subcloned from cosmid 166G8 into pBluescriptII (SK) (Stratagene). The clone was sequenced using prim-ers derived from the bPABII gene and human EST sequences. Sequencing of the PAB II introns was done by primer walking.

PAB II mutation screening and sequencing

All cases were diagnosed as having OPMD on clinical grounds (Brais, B. et al., Hum. Mol. Genet. 4, 429-434 (1995)). RT-PCR- and genomic SSCP analyses were done using stan-dard protocols (Lafrenière, R.G. et al., Nat. Genet. 15, 298-302 (1997)). The primers used to amplify the PAB II mutated region were: 5'-CGCAGTGCCCCGCCTTAGA-3' (SEQ ID NO:4) and 5'-ACAAGATGGCGCCGCCGCCCGGC-3' (SEQ ID NO:5). PCR reactions were performed in a total volume of 15 µl containing: 40 ng of genomic DNA; 1.5 µg of BSA; 1 µM of each primer; 250 µM dCTP and dTTP; 25 µM dATP; 125 µM of dGTP and 125 µM of 7-deaza-dGTP (Pharmacia); 7.5% DMSO; 3.75 µCi[³⁵S]dATP, 1.5 unit of Taq DNA polymerase and 1.5 mM MgCl₂ (Perkin Elmer). For non-radioactive PCR reactions the [³⁵S]dATP was replaced by 225 µM of dATP. The amplification procedure consisted of an initial denaturation step at 95°C for five minutes, followed by 35 cycles of denaturation at 95°C for 15 s, annealing at 70°C for 30 s, elongation at 74°C for 30 s and a final elongation at 74°C for 7 min. Samples were loaded on 5% polyacrylamide denatur-ing gels. Following electrophoresis, gels were dried and autoradiographs were obtained. Sizes of the inserts were determined by comparing to a standard M13 sequence (Sequenase™, USB). Fragments used for sequencing were gel-purified. Sequencing of the mutated fragment using the Amplicycle kit™ (Perkin Elmer) was

done with the 5'-CGCAGTGCCCCGCCTTAGAGGTG-3' (SEQ ID NO:6) primer at an elongation temperature of 68°C.

Stability of (GCG)-repeat expansions

The meiotic stability of the (GCG)₉-repeat was estimated based on a large French Canadian OPMD cohort. It had been previously established that a single ancestral OPMD carrier chromosome was introduced in the French Canadian population by three sisters in 1648. Seventy of the seventy one French Canadian OPMD families tested to date segregate a (GCG)₉ expansion. However, in family F151, the affected brother and sister, despite sharing the French Canadian ancestral haplotype, carry a (GCG)₁₂ expansion, twice the size of the ancestral (GCG)₉ mutation (Fig. 2C). In this founder effect study, it is estimated that 450 (304-594) historical meioses shaped the 123 OPMD cases belonging to 42 of the 71 enrolled families. The screening of the full set of participants allowed an identification of another 148 (GCG)₉ carrier chromosomes. Therefore, it is estimated that a single mutation of the (GCG)₉ expansion has occurred in 598 (452-742) meioses.

Genotype-phenotype correlations

176 carriers of at least one copy of the (GCG)₉ mutation were examined during the early stage of the linkage study. All were asked to swallow 80 cc of ice-cold water as rapidly as possible. Testing was stopped after 60 seconds. The swallowing time (st) was validated as a sensitive test to identify OPMD cases (Brais, B. et al., Hum. Mol. Genet. 4, 429-434 (1995); Bouchard, J.-P. et al., Can. J. Neurol. Sci. 19, 296-297 (1992)). The st values for 76 (GCG)₆ homozygotes normal controls is illustrated in Fig. 3. Analyses of variance were computed by two-way ANOVA (SYSTAT package). For the (GCG)₉ homozygotes their mean st value was compared to the mean value for all (GCG)₉ heterozygotes aged 35-40 ($P < 10^{-5}$). For the (GCG)₉ and (GCG)₇ compound

91

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth, and as follows in the scope of the appended claims.

CLAIMS

What is claimed is:

1. An isolated human PAB II gene comprising a polymorphic GCG repeat in exon I thereof wherein an allelic variant of said GCG repeat is indicative of a disease associated with protein accumulation in a cell nucleus, swallowing difficulty, and/or ptosis in a human patient.
2. The gene of claim 1, wherein said polymorphic GCG repeat has the sequence
ATG (GCG)_{6+n} GCA,
wherein n is selected from 1 to 7.
3. The gene of claim 2, wherein n is selected from 2 to 7, and wherein said allelic variant is associated with an increased severity of the disease.
4. The gene of claim 3, wherein a phenotype associated with said allelic variant is dominant.
5. The gene of claim 2, wherein a first allele of said GCG repeat has an n which is equal to 1.
6. The gene of claim 5, wherein a second allele of said GCG repeat has an n selected from 2 to 7, and wherein said first allele is a modulator of the severity of the phenotype associated with said second allele.
7. The gene of claim 1, wherein said human patient is homozygous for said polymorphic GCG repeat.

8. The gene of claim 1, wherein said human patient is heterozygous for said polymorphic GCG repeat.
9. A nucleic acid sequence comprising a polymorphic GCG repeat of exon I of a human PAB II gene, wherein an allelic variant of said GCG repeat in a patient's human PAB II gene is indicative of a disease associated with protein accumulation in a cell nucleus, swallowing difficulty, and/or ptosis in said human patient.
10. The nucleic acid sequence of claim 9, wherein said polymorphic GCG repeat has the sequence

$$\text{ATG (GCG)}_{6+n} \text{GCA},$$
wherein n is selected from 1 to 7.
11. The nucleic acid sequence of claim 10, wherein n is selected from 2 to 7, and wherein said allelic variant is associated with an increased severity of said disease.
12. The nucleic acid sequence of claim 11, wherein a phenotype associated with said allelic variant is dominant.
13. A method for the diagnosis or prognosis of a disease associated with protein accumulation in a cell nucleus, and/or swallowing difficulty and/or ptosis in a human patient, which comprises:
 - a) obtaining a nucleic acid sample of said patient; and
 - b) determining allelic variants of a GCG repeat in exon I of the PAB II gene, said GCG repeat having the sequence



wherein n is selected from 0 to 7, and
whereby at least one allele of said GCG repeat having an n equal to 1 to 7, is
indicative of a disease related with said protein accumulation in said nucleus,
and/or a swallowing difficulty and/or ptosis in said patient.

- 5 14. The method of claim 13, wherein n is selected from 2 to 7, and wherein said
 allelic variant is associated with an increased severity of said disease.
15. The method of claim 14, wherein said phenotype associated with said allelic
 variant is dominant.
- 10 16. The method of claim 13, wherein a first allele of said GCG repeat has an n
 which is equal to 1.
17. The method of claim 16, wherein a second allele of said GCG repeat has an n
 selected from 2 to 7, and wherein said first allele is a modulator of the severity
 of the phenotype associated with said second allele.
18. The method of claim 13, wherein said disease is oculopharyngeal muscular
15 dystrophy.
19. A non-human transgenic animal whose germ cells and somatic cells are
 modified to express at least one allelic variant of a polymorphic GCG repeat in
 exon I of the PAB II gene, and wherein said transgenic animal shows a
 phenotype of a disease associated with protein accumulation in a cell nucleus,
20 and/or a swallowing difficulty and/or ptosis.

20. The non-human transgenic animal of claim 19, wherein said polymorphic GCG repeat has the sequence
- $$\text{ATG (GCG)}_{6+n} \text{GCA,}$$
- wherein n is selected from 1 to 7.
- 5 21. The transgenic animal of claim 19, wherein said animal is a mammal.
22. The transgenic animal of claim 19, wherein said allelic variant of the PAB II gene is a human allelic variant.
23. The transgenic animal of claim 19, having cells which display a protein accumulation in their nucleus.
- 10 24. A cell isolated from said non-human transgenic animal according to claim 19.
25. A method for screening and identifying an agent for the prevention and/or treatment of a disease associated with protein accumulation in a cell nucleus and/or a swallowing difficulty, and/or ptosis, said method comprising:
- 15 a) exposing the transgenic animal of claim 19 to said agent; and
- b) evaluating the prevention and/or treatment of development of said protein accumulation in a cell nucleus and/or a swallowing difficulty, and/or ptosis in said animal exposed to said agent as compared to a control animal not having been exposed to said agent.
26. A method for screening and identifying an agent which modulates protein accumulation in the nucleus of a cell, said method comprising:
- 20 a) exposing a cell of claim 24 to said agent; and

- b) evaluating said protein accumulation in said nucleus of said exposed cell as compared to a control cell not having been exposed to said agent.
27. The method of claim 25, wherein said protein accumulation is associated with oculopharyngeal muscular dystrophy.
- 5 28. A cell which has been modified to express at least one allelic variant of a human polymorphic GCG repeat of exon I of the PAB II gene, wherein said allelic variant is associated with protein accumulation in the nucleus of said cell.
29. The cell of claim 28, wherein said polymorphic GCG repeat has the sequence
ATG (GCG)_{6+n} GCA,
10 wherein n is selected from 1 to 7.
30. The cell of claim 28, wherein said cell is a mammalian cell.
31. An isolated human PAB II gene comprising a polymorphic GCG repeat in exon I thereof, wherein said repeat has the sequence ATG (GCG)_{6+n} GCA, wherein n is 0, and wherein said sequence is indicative of a non-disease phenotype
15 associated with protein accumulation in a cell nucleus, swallowing difficulty, and/or ptosis in a human patient.
32. The human PAB II gene of claim 31, wherein said gene is as set forth in SEQ ID NO:3.
33. A method of diagnosing a disease in a human patient associated with a
20 meiotically stable trinucleotide expansion in a coding sequence of a gene comprising:

- 5

10

36. The method of claim 33, wherein said disease is oculopharyngeal muscular dystrophy.

ABSTRACT OF THE DISCLOSURE

The present invention relates to a human PAB II gene containing transcribed polymorphic GCG repeat, which comprises a sequence as set forth in SEQ ID NO:3, which includes introns and flanking genomic sequence. The allelic variants of GCG repeat of the human PAB II gene are associated with a disease related with protein accumulation in nucleus, such as polyalanine accumulation, a disease related with swallowing difficulties, such as oculopharyngeal muscular dystrophy. The present invention also relates to a method for the diagnosis of a disease with protein accumulation in nucleus, which comprises the steps of: a) obtaining a nucleic acid sample of said patient; and b) determining allelic variants of GCG repeat of the gene of claim 1, and wherein long allelic variants are indicative of a disease related with protein accumulation in nucleus.




FIG. 1A

Human *PAB II* genomic structure (6002 bp)




FIG. 1B




FIG. 3

09590211.060800

1 aatgaagtg gacacccaat tagccccaat acaatgctt gttcaatcaa ccaaacatct
61 aagcagcaca tctatgtgt agcatatgc caguccgtga gactgcgaat ataatagga
121 accgcccctc atctgcagc gctcacaacc tagttaagcaa acagtaaac attaagcgc
181 gccgtggaca taggcccaat tgtcctggga aatgaggga agctgggtt tgcagtggt
241 tgattgaag gggaactacat gttagaggca cagactgggt gcaggtacac ccaaggaac
301 gagaagagt gaagaaaaca acatccaaca agtaaccaca tgctgcgta tcgaaggccg
361 tgatttacg tttgagact ttacctgcc agcaaaaggg gccagttcty ttagcgtgc
421 agattgagg gtgacattg gaagctgtcc aguaaaaaga aaatggaact gggagcaga
481 aggcctacgc aagaggcgg gacagacag acttgtagt agtagtcty gactgagga
541 tcctccctgc ttctggtgc gggagagcta gtgatgatg gtccaataa cctgatgg
601 gaaagtaagc tccctcctg aatgcttcat tcacaacctc catttcagc aacatccat
661 ctactggtgc ttcctggtcg agatacaagt ttcctgaaac tgctgctcty ttltggcct
721 cacccgcca acagctcaat agctggcaag cagtagtatc aagatggcgg cccctagga
781 ctggtagtc atgtgaacct ggtttccca agtttgaagc ccggcagtc ttctggggc
841 aaggttcacc tgtcacgaaa cgagtgtcac cccttcgact ctgcaagcc aatcggcatc
901 tgagactgg ccactgcggt gaggcgatcg gaagatlggt ccttccagt cgcctagcta
961 ggccaatca cggagcgtcc catacttcgc gggcccggcc gtaggccgg gagaaucag
1021 aatatcgtca cagcgttgcg gtattattac gtaagactc ctaagactc gataggagt ggagcgtg
1081 ttgattgaca ggcagatttc cctaccggga tttagaatc ttggcgagt cccgccttag
1141 aggtgcgctt atttgattgc caagtaatat tcccgaatgg agtactagct catggtgacg

FIG. 4A

1201 ggcagggcagc ttgagctaat gagtccctccg ttggccggcgcg agctctccac atgccggggcg
1261 gccgggccccca gtcttgagcgg cgatggcggc gccgggcggcg gccggcagcag caggcggggcg
1321 tgcggggcgggt cggggctccg ggccggggggcg gccggcgccat ctgtgtgccg gggccgggtg
1381 ggagggccgggg gagggggccc cggggggcgcg aggggactac gggaacggcc tggagtctga
1441 ggaactggag cctgaggagc tgctgtgga gcccgagccg gagcccgagc ccgaaggagga
1501 gccgcccccg ccccgcgccc ccccgggagc tccgggccct gggtctgtt cgggagcccc
1561 cggcagccaa gaggaaggag aggaagccggg actgtcga ggtgaaccgg gggacggcgcg
1621 cattgagga cccgtgagga aggaaggcgga gccgaagcagc cggcggtgg cgcgtcactg
1681 gagggcccaga gctcggggcga gccgtggcag gccgggggtg gggtlgygc gggaataacg
1741 tggctggggc gggtcgggcc gggtatgggt cagcgatcac tacaaggggc ccgaactggct
1801 tgattcgggc gtcacgggtg cctagtgtg ttctaaggag ggtagcttt ctttatcac
1861 gacctcgca ttggggcga gaaatggccg agcatggctg aggcggctc ttggccgagag
1921 cagggcacag cccctgcgtt ggttcctct aagctgtcct ccataccctc cccactata
1981 ttaggagctg gaagctatca aagctcgagt cagggagatg gaggaagaa ctgagaagct
2041 aaaggagcta cagaaacgag tagaagaagca gatgaatatg agtccaacc caggcaatgc
2101 tgaagtaactg gcggttgca cgcgagcccg ggttctcggg ttggaagggt tgtggggagg
2161 atggggaatg ttgggttaqa tactcggcac cctggagctg ctgtctgag ctattatgac
2221 tgtgccggcg tcatagtccg ttgtgtgttc ctctgacct ttgtaggcag aactgatatt
2281 ttgtgtgtg tagcccttg cctcccttg tcctgttata attgtgttg tctttattct
2341 tagtctaagt ctatcttct ttgttagagg ttgcgtgctc gcaattgacc ttcaaatcta

FIG. 4B

2401 atagtttttc ctccaattgg agacgcctta ggaattcctaag agaaagcaag ctggaagggy
2461 ttccccctt aatltctaga aatgtgagt ctcaagcccac ttaattttgc tcaactcta
2521 aagcatttca accaaagcca ttcattaggg atttgatttg gagggcagga gggattccta
2581 tactgttta agtgtgtatt aattccttca attatcgaa ttatttagtg agtaaccctgc
2641 tatgcactag gcactattct cggcttgtgg gtacagcaggy gaacagcacca gaccaaaatc
2701 ttgaccttca ctgagcctat gggatagtcg tgytgytgga agtgcaacat attggtcaag
2761 tagaaaaacaa gtgtgtggt ttgttaaaaa attatlttt cctgatatgt gggccggtga
2821 tcatgtccat tgagaggaag atggaagctg atgcccgctc catctatgtt ggcaatgtga
2881 cgtactgggg ctctgactgg ggttgggggc aagttccttc ttgggggaat tatttaatag
2941 tcctgaaaga acatctccgg gatagatgtg gtttgggtg tggagggagt gtgggaagga
3001 ggttaaaagt aatggaatga tcaagtaatca gcaaaaggctc tgggtttgga agyaaagag
3061 attaatccct caaatlaccg gatttcatgt gctttggtgt atgatgccc agaccaaagg
3121 ctcgggaggg ttcttttgag acaggaattt gcctggtgcc tgtgaaattt ttctcctc
3181 atcaggttga ctatgttga acagcagaag agctggaagc tcactttcat ggctgttgtt
3241 cagtcaaccg tgtlaccata ctgttgaca aatttagtgg ccatcccaa ggtaaagtaa
3301 agggagtaaa gttgagataa tttaatttac agtgtlaca aa atgtttata
3361 ttgagcagta agttatttg tgttaacaca ggtgatctgt gtcatttaag atcatggcat
3421 taatgtttgat atatcaggag ttgcaacctaa atgtcctcag aggccagata acaaaatga
3481 aggcatagat tgggtgggat tacgaaactag aaggggaggg gcagcttcta ctggcctat
3541 tatggcatat ggaaatlcag gccctgtgtg tcttattttt acaaatltca aagagtagct

FIG. 4C

3601 ggaatttta aaatttaaat gatttcgaat gattgaaatt ttccatttag aagaattttg
3661 acaataaaaa aatatlaactg catgttagcc caaaacgaaag catgcctgca ggttgaattt
3721 gaacctgtgag gtatttghta cctcagagag atacaatgac aattccttc aggtttgct
3781 atatagagtt ctcaagacaaa gagtcagtgga ggaacttcctt ggccttagat ggttccctat
3841 ttagaggaag gcaaatcaag gtaagcctat gtccattgct gttctagttg tghtataact
3901 ctccaggttg cctttaagc tatcatttgt tcatctctga ctcaagtgat cccaaaacga
3961 accaacagac caggcattcag caccaacagac cggggtttc caccagcccg ctaccgcgcc
4021 cgaaccacca actacaacag ctcccgctc cgaatctaca gtggtltaa cagcagggcc
4081 cggggtcgcg tctacagtc aggatagatg ggctgctcct cttcccccg cctcccgtag
4141 gccccgtatg cttcctcctc tctggtctga ggaacctccc tcccccacc cctcccgtag
4201 gtcttcagga acttgtctc ctgcctgtgc aggttgagga aggtagttgc aggcagggcc
4261 agaaggcagc ctcatcatct ttctgcagt agaatgtgt gataaggct gcatccctcc
4321 ctgtgtcaa agaggttcc accccagcc ttttttctt tgggagttgg tgcatltag
4381 agtgtttgc ggaacaaact ggagggaaac gggcctccag gaagltgaaa gcactgctg
4441 gacatttgtt acttttttcg gaattagga gggatlgaa actgaaacct ccttggaaga
4501 ataccagag ctagctagt gatcctcca acagcctgtt ggagagatt tgagataact
4561 attctttat tgagccagtc ttgcaagtt aactctcac tgggcctagt gtgtncca
4621 ggtttttgcc ttgcttcaact tctgtctcta catttaata gacgggttag gcatataaac
4681 ctggtcttt cataagctct acctgcctat cccagggagt tagggaggt ctatlytgaa
4741 aggccctagg gtttaaaac tgtgagggac tgaaaaactg gataaaaagg ggtccttlt

FIG. 4D

```

4801 ccttgccctt gtctctcact cagatgcgct tcttttcgc cactgttttg caaagtttc
4861 tghtaagccc ccctccccc gccccaagttc tcccaagttgc gttactattt ctgggatcat
4921 ggggtcgggt ttagaacact tgaacacttc ttccccccc ttcccttcac agtaactggg
4981 gcaggggcct acggggaggg gcttgtactg aactatctag tgatcacggt aaccactaac
5041 tctccttctt tcttcaggg gccgggctag agcggacatca tgytaltccc cttaactaaa
5101 aaagtgtgtt ttaggaggag agagaggaaa aaaaaggaaa agaaagaaaa tttttaaa
5161 taaaaaaaaa aaaaaagaaa acagaagatg acctgatgg aaaaaaata tttttaaa
5221 aaaaatatata ctgtggaag ggggagaatc ccataactaa ctgctgagg gggacctgct
5281 ttggggagta ggggaaagcc cagggagttg ggcagggggc tgcttattca ctctgggat
5341 tcgccatgga cagttctcaa ctgcgcaagc tgcttgccca tgtttccctg ccccttcac
5401 ccccttggc ctgtcaagg gtaggtggc gtgggtgtta ggaggttt tttaaccag
5461 ggctctgga ggaacaccaa ctgttctgct tgttacctc cctccgctc tctcctgcc
5521 ttcacagtc ccctcctgcc tgcctctgtc cagccaggtc taccacccac ctttgtgtt
5581 ttctccggc tccctgcccc tccagattgc ctgtgatct atttgttc ctttgtgtt
5641 tcttttctg ttttagtgt ctctcttgcc aggtttctgt agccggaaga tctccgttc
5701 gctcccaagg gctccagtgt aaattcccc tccccctggg gaaatgcact acctgtttt
5761 ggggggttta ggggtgttt ttgttttcag ttgttttgt ttttgttt ttltttcc
5821 tttagctttt ttccctttta ttggaaggaa atgggaaggaa gtgggaacag ggaagtggga
5881 ggtgatttt ttatttttt ttagctcatt tccagggtg ggaattttt tttaatatgt
5941 gtcatgaata aagttgttt ttgaaaataa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa
6001 aa

```

FIG. 4E

SEQUENCE LISTING

<110> MCGILL UNIVERSITY
ROULEAU, Guy A.
BRAIS, Bernard

<120> SHORT GCG EXPANSIONS IN THE PAG II GENE
FOR OCULOPHARYNGEAL MUSCULAR DYSTROPHY AND DIAGNOSTIC
THEREOF

<130> 3028.1000-000

<150> CA 2,218,199

<151> 1997-12-09

<160> 6

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 57

<212> DNA

<213> Artificial Sequence

<400> 1

atggcggcgg cggcggcgcg ggcagcagca atggcggcgg cggcggcgcg ggcggca 57

<210> 2

<211> 35

<212> DNA

<213> Artificial Sequence

<400> 2

maaaaaaaaa agaaggrcsm aaaaaaaaaa agaag 35

<210> 3

<211> 6002

<212> DNA

<213> Artificial Sequence

<400> 3

aatgaaggtg	gacaccca	tagcccca	acaaatgc	gttcaatcaa	ccaaacatct	60
aagcagcaca	tctatgtggt	agcatattgc	caggccgtga	gactgcgaat	ataaatagga	120
accgcccctc	atctgcaggc	gctcacaacc	tagttagcaa	acagtaaaac	aattaagcgc	180
gccgtggaca	taggcccact	tgctctggga	aatgagggga	agctgggggt	tgcaagtgggt	240
tgattgaagg	gggactacat	gttagaggca	cagactgggt	gcaggtacac	ccaaagggaac	300
gagaagagtg	gaaggaaaca	acatccacaa	agtaaccaca	tgctggcgta	tcgaaggccg	360
tgatttacgg	ttttgagact	ttacctcgcc	agcaaagggg	ggccagtctg	ttagcgggtg	420
agattggagg	ggtgacattg	gaagctgtcc	aggaaaaaga	aaatggaact	ggggagcaga	480
aggcctacgc	aagagggcgg	gacagacagg	acttgtgact	agtagctctg	gactgaggaa	540
tcctccctgc	tttctgggtg	gggagagcta	gtggatgatg	gtgccaataa	cctggatggg	600
gaaagtaagc	tcctcctcgg	aatgcttcat	tcacaacctc	cattttcagc	aacatcccat	660
ctactggtgc	ttctgggtcg	agatacaagt	ttctgaaac	tgctgctctg	ttttgggcct	720
caccgcggcca	acagctcact	agctggcaag	cagtagtatc	aagatggcgg	ccccctagga	780
ctggctagtc	atgtgacctc	gggtttccca	agtttgaagc	ccggcagtc	tttcgggggc	840
aaggttcacc	tgtcacgaaa	cgagtgtcac	cccttcgact	ctcgcaagcc	aatcggcatc	900
tgagactggg	ccactgcggt	gaggcgatcg	gaagattggg	cctttccagt	cgcctagcta	960
gggccaatca	cggagcgtcc	catacttcgc	gggcccgcgc	gtaggccggg	gagaagcagg	1020


```
<210> 4
<211> 19
<212> DNA
<213> Artificial Sequence
```

```
<210> 5
<211> 24
<212> DNA
<213> Artificial Sequence
```

```
<210> 6
<211> 23
<212> DNA
<213> Artificial Sequence
```

